arXiv:1105.4307v2 [math.RA] 9 Apr 2012

Algebra with Conjugation

Aleks Kleyn

Abstract. In the paper, I consider properties and mappings of free algebra with unit. I consider also conjugation of free algebra with unit.

Contents

1. Preface . . . 1
2. Conventions . . . 2
3. Algebra with unit . . . 2
4. Algebra with Conjugation . . . 4
5. Antilinear Mapping of Algebra with Conjugation . . . 8
6. References . . . 9
7. Index . . . 10
8. Special Symbols and Notations . . . 11

1. Preface

When I started to write the book [2], I realized that classical definition of a linear mapping

$$y = ax \quad y = xa$$

restricts our possibilities to study the derivative of mapping into division ring. Intuitively, it was clear that we can represent the linear mapping into division ring as a sum of terms like

$$axb$$

Following study of the structure of linear mapping into division ring was finished in the book [4]. I expanded the circle of considered algebras and started to study free algebra over commutative ring.

If the set A is D-algebra, then we can define few algebraic structures on the set A. According to the definition [4]-2.2.1, A is D-module. If we consider only sum and product in D-algebra A, then we can consider D-algebra A as ring (as option, the ring maybe nonassociative). According to this, we can consider D-algebra A as $A\star$-module of dimension 1. This is why it is important to study mappings that preserve one or another algebraic structure on the set A.

In this paper, I considered $A\star$-linear mapping of $D\star$-algebra A.

Aleks_Kleyn@MailAPS.org.
`http://sites.google.com/site/AleksKleyn/`.
`http://arxiv.org/a/kleyn_a_1`.
`http://AleksKleyn.blogspot.com/`.

The concept of algebra with conjugation fits naturally into the theory of D-algebra with unit. However I believe that we can define conjugation even in case if algebra does not have unit. However the answer on this question requires further research.

2. CONVENTIONS

Convention 2.1. *Let A be free finite dimensional algebra. Considering expansion of element of algebra A relative basis $\overline{\overline{e}}$ we use the same root letter to denote this element and its coordinates. However we do not use vector notation in algebra. In expression a^2, it is not clear whether this is component of expansion of element a relative basis, or this is operation $a^2 = aa$. To make text clearer we use separate color for index of element of algebra. For instance,*

$$a = a^{\boldsymbol{i}}\overline{e}_{\boldsymbol{i}}$$

$\square$

Convention 2.2. *If free finite dimensional algebra has unit, then we identify the vector of basis $\overline{e}_{\mathbf{0}}$ with unit of algebra.* $\square$

Convention 2.3. *Although the algebra is a free module over some ring, we do not use the vector notation to write elements of algebra. In the case when I consider the matrix of coordinates of element of algebra, I will use vector notation to write corresponding element. In order to avoid ambiguity when I use conjugation, I denote a^* element conjugated to element a.* $\square$

Without a doubt, the reader may have questions, comments, objections. I will appreciate any response.

3. ALGEBRA WITH UNIT

Let D be commutative ring. Let A be D-algebra with unit 1. Therefore there exists effective representation

$$f_{1,2} : d \circ a = da \quad d \in D \quad a \in A \tag{3.1}$$

of the ring D in the algebra A.

Theorem 3.1. *Let D be commutative ring. Let A be D-algebra. Let*

$$f : A^n \to A$$

be n-linear mapping into D-algebra A. Let

$$g : A^m \to A$$

be m-linear mapping into D-algebra A. Then if for given i in the expression

$$f \circ (x_1, ..., x_n)$$

we make the substitution

$$x_i = g \circ (y_1, ..., y_m)$$

then the resulting expression

$$h(x_1, ..., \widehat{x_i}, ..., x_n, y_1, ..., y_m) = f \circ (x_1, ..., g \circ (y_1, ..., y_m), ..., x_n)$$

is $(m + n - 1)$-linear mapping.

Proof. Since polylinear mapping is linear mapping with respect to each argument, then the mapping h is linear with respect to x^j, $j \neq i$. Since the mapping f is linear mapping with respect to x^i and for each k the mapping g is linear mapping with respect to y^k, then according to the theorem [4]-2.4.1, the mapping h is linear mapping with respect to y^k. □

Theorem 3.2. *Let D be commutative ring. Let A be D-algebra. Then*

$$d(ab) = (da)b = a(db) \quad d \in D \quad a, b \in A \tag{3.2}$$

Proof. Let f be bilinear mapping corresponding to product in D-algebra A.[1] Then

$$df(ab) = f(da, b) = f(a, db) \quad d \in D \quad a, b \in A \tag{3.3}$$

The equation (3.2) follows from the equation (3.3). □

Theorem 3.3. *Let D be commutative ring.*[2] *The commutator of D-algebra A is bilinear mapping. The associator of D-algebra A is 3-linear mapping.*

Proof. According to definitions [4]-2.2.1, [4]-2.2.4, the commutator is bilinear mapping. According to definitions [4]-2.2.1, [4]-2.2.5 and the theorem 3.1, the associator is 3-linear mapping. □

Theorem 3.4. *Let D be commutative ring. Let A be D-algebra with unit e. Then we can identify $d \in D$ and $d\,1 \in A$. In such case, $D \subseteq Z(A)$.*[3]

Proof. From the theorem 3.3, it follows that

$$\begin{aligned}
(d, a, b) &= d(1, a, b) = d((1a)b - 1(ab)) = d(ab - ab) = 0\\
(a, d, b) &= d(a, 1, b) = d((a1)b - a(1b)) = d(ab - ab) = 0\\
(a, b, d) &= d(a, b, 1) = d((ab)1 - a(b1)) = d(ab - ab) = 0
\end{aligned}$$

Therefore, $D \subseteq N(A)$. If we assume $b = 1$ in the equation (3.2), then we get

$$da = ad \quad d \in D \quad a \in A \tag{3.4}$$

From the equation (3.4), it follows that the commutator has form

$$[d, a] = 0$$

□

Theorem 3.5. *Structural constants of D-algebra with unit e satisfy condition*

$$C^{l}_{0k} = C^{l}_{k0} = \delta^{k}_{l} \tag{3.5}$$

Proof. The equation (3.5) follows from the equation $1d = d1 = d$. □

---

[1] I consider the definition of the product in D-algebra according to the definition [4]-2.2.1.

[2] You can see the definition of commutator in the definition [4]-2.2.4. You can see the definition of associator in the definition [4]-2.2.5. You can see also [5], p. 13.

[3] You can see the definition of nucleus $N(A)$ of algebra A in the definition [4]-2.2.7; you can see the definition of center $Z(A)$ of algebra A in the definition [4]-2.2.8; see also [5], p. 13, 14.

Theorem 3.6. *$A\star$-linear mapping of D-algebra A with unit has form*

$$f(c)=cd\quad d\in A \tag{3.6}$$

Let $\overline{\overline{e}}$ be $(\mathrm{Re}\,A)$-basis of the algebra A. Coordinates of the mapping f

$$f(c)=c^l f^i_l\,\overline{e}_i \tag{3.7}$$

relative to the basis $\overline{\overline{e}}$ satisfy to the equation

$$f^j_k=f^i_0C^j_{ki} \tag{3.8}$$

$$d=f^i_0\,\overline{e}_i \tag{3.9}$$

Proof. $A\star$-linear mapping satisfies to equations

$$f(ac)=(ac)^i f^j_i\overline{e}_j=a^kc^lC^i_{kl}f^j_i\overline{e}_j \tag{3.10}$$

$$af(c)=a^k(f(c))^iC^j_{ki}\overline{e}_j=a^kc^lf^i_lC^j_{ki}\overline{e}_j \tag{3.11}$$

From equations (3.10), (3.11), it follows that

$$a^kc^lC^i_{kl}f^j_i\overline{e}_j=a^kc^lf^i_lC^j_{ki}\overline{e}_j \tag{3.12}$$

The equation

$$C^i_{kl}f^j_i=f^i_lC^j_{ki} \tag{3.13}$$

follows from the equation (3.12).

From the equation (3.13) it follows that[4]

$$C^i_{k0}f^j_i=f^i_0C^j_{ki} \tag{3.14}$$

From equations (3.5), (3.14), it follows that

$$\delta^i_kf^j_i=f^i_0C^j_{ki} \tag{3.15}$$

The equation (3.8) follows from the equation (3.15).

From equations (3.7), (3.8), it follows that

$$f(c)=c^lf^j_0C^i_{lj}\overline{e}_i \tag{3.16}$$

The equation (3.9) follows from the equations (3.6), (3.16). □

4. ALGEBRA WITH CONJUGATION

Let D be commutative ring. Let A be D-algebra with unit e, $A\neq D$. According to the theorem 3.4, we can identify the ring D and subalgebra of D-algebra A. Therefore, D-algebra A has a nontrivial center $Z(A)$.

Let there exist subalgebra F of algebra A such that $F\neq A$, $D\subseteq F\subseteq Z(A)$, and algebra A is a free module over the ring F. Let $\overline{\overline{e}}$ be the basis of free module A over ring F. We assume that $\overline{e}_0=1$.

Consider mappings

$$\mathrm{Re}:A\rightarrow A$$
$$\mathrm{Im}:A\rightarrow A$$

defined by equation

$$\mathrm{Re}\,d=d^0\quad \mathrm{Im}\,d=d-d^0\quad d\in D\quad d=d^i\overline{e}_i \tag{4.1}$$

---

[4] Let $l=0$.

The expression $\mathrm{Re}\, d$ is called **scalar of element** d. The expression $\mathrm{Im}\, d$ is called **vector of element** d.[5]

According to (4.1)

$$F = \{d \in A : \mathrm{Re}\, d = d\}$$

We will use notation $\mathrm{Re}\, A$ to denote **scalar algebra of algebra** A.

Theorem 4.1. *The set*

$$\mathrm{Im}\, A = \{d \in A : \mathrm{Re}\, d = 0\} \tag{4.2}$$

is $(\mathrm{Re}\, A)$*-module which is called* **vector module of algebra** A*.*

$$A = \mathrm{Re}\, A \oplus \mathrm{Im}\, A \tag{4.3}$$

Proof. Let c, $d \in \mathrm{Im}\, A$. Then $c_0 = d_0 = 0$. Therefore,

$$(c+d)_0 = c_0 + d_0 = 0$$

If $a \in \mathrm{Re}\, A$, then

$$(ad)_0 = ad_0 = 0$$

Therefore, $\mathrm{Im}\, A$ is $(\mathrm{Re}\, A)$-module.

Sequence of modules

$$\begin{tikzcd} 0 \arrow[r] & \mathrm{Re}\, A \arrow[r, "\mathrm{id}"] & A \arrow[r, "\mathrm{Im}"] & \mathrm{Im}\, A \arrow[r] & 0 \end{tikzcd}$$

is exact sequence. According to the definition (4.1) of the mapping Re, following diagram is commutative

$$\begin{tikzcd} \mathrm{Re}\, A \arrow[r, "\mathrm{id}"] \arrow[d, "\mathrm{id}"'] & A \arrow[ld, "\mathrm{Re}"] \\ \mathrm{Re}\, A & \end{tikzcd}$$

According to the statement 2 of the proposition [1]-III.3.2,

$$A = \mathrm{id}(\mathrm{Re}\, A) \oplus \ker \mathrm{Re} \tag{4.4}$$

According to the definition (4.2)

$$\ker \mathrm{Re} = \{d \in A : \mathrm{Re}\, d = 0\} = \mathrm{Im}\, A \tag{4.5}$$

The equation (4.3) follows from the equations (4.4), (4.5). □

According to the theorem 4.1, there is unique defined representation

$$d = \mathrm{Re}\, d + \mathrm{Im}\, d \tag{4.6}$$

Definition 4.2. The mapping

$$d^* = \mathrm{Re}\, d - \mathrm{Im}\, d \tag{4.7}$$

is called **conjugation in algebra** provided that this mapping satisfies

$$(cd)^* = d^*\, c^* \tag{4.8}$$

$(\mathrm{Re}\, A)$-algebra A equipped with conjugation is called **algebra with conjugation**. □

Corollary 4.3. $(d^*)^* = d$ □

---

[5] In the section [3]-7, I consider an example of algebra in which there are two conjugation. We assume that we chose conjugation.

Theorem 4.4.

$$d+d^*\in\operatorname{Re}A \tag{4.9}$$

$$d-d^*\in\operatorname{Im}A \tag{4.10}$$

Proof. The theorem follows from equations (4.6), (4.7). □

Theorem 4.5. *The* $(\operatorname{Re}A)$*-algebra* A *is algebra with conjugation if structural constants of* $(\operatorname{Re}A)$*-algebra* A *satisfy condition*

$$C^{\mathbf{0}}_{\boldsymbol{kl}}=C^{\mathbf{0}}_{\boldsymbol{lk}}\qquad C^{\boldsymbol{p}}_{\boldsymbol{kl}}=-C^{\boldsymbol{p}}_{\boldsymbol{lk}} \tag{4.11}$$

$$\mathbf{1}<\boldsymbol{k}<\boldsymbol{n}\quad \mathbf{1}<\boldsymbol{l}<\boldsymbol{n}\quad \mathbf{1}<\boldsymbol{p}<\boldsymbol{n}$$

Proof. From equations (4.6), (4.7) it follows that

$$\begin{aligned}(cd)^*&=(\operatorname{Re}c\ \operatorname{Re}d+\operatorname{Re}c\ \operatorname{Im}d+\operatorname{Im}c\ \operatorname{Re}d+\operatorname{Im}c\ \operatorname{Im}d)^*\\&=\operatorname{Re}c\ \operatorname{Re}d-\operatorname{Re}c\ \operatorname{Im}d-\operatorname{Im}c\ \operatorname{Re}d+(\operatorname{Im}c\ \operatorname{Im}d)^*\end{aligned} \tag{4.12}$$

$$\begin{aligned}d^*\,c^*&=(\operatorname{Re}d-\operatorname{Im}d)(\operatorname{Re}c-\operatorname{Im}c)\\&=\operatorname{Re}d\ \operatorname{Re}c-\operatorname{Re}d\ \operatorname{Im}c-\operatorname{Im}d\ \operatorname{Re}c+\operatorname{Im}d\ \operatorname{Im}c\end{aligned} \tag{4.13}$$

From equations (4.8), (4.12), (4.13), it follows that

$$(\operatorname{Im}c\ \operatorname{Im}d)^*=\operatorname{Im}d\ \operatorname{Im}c \tag{4.14}$$

Let $\overline{\overline{e}}$ be the basis of the $(\operatorname{Re}D)$-algebra D. From the equation (4.14), it follows that

$$\begin{aligned}(C^{\mathbf{0}}_{\boldsymbol{kl}}c^{\boldsymbol{k}}d^{\boldsymbol{l}}+C^{\boldsymbol{p}}_{\boldsymbol{kl}}c^{\boldsymbol{k}}d^{\boldsymbol{l}}\overline{e}_{\boldsymbol{p}})^*&=C^{\mathbf{0}}_{\boldsymbol{kl}}c^{\boldsymbol{k}}d^{\boldsymbol{l}}-C^{\boldsymbol{p}}_{\boldsymbol{kl}}c^{\boldsymbol{k}}d^{\boldsymbol{l}}\overline{e}_{\boldsymbol{p}}\\&=C^{\mathbf{0}}_{\boldsymbol{kl}}d^{\boldsymbol{k}}c^{\boldsymbol{l}}+C^{\boldsymbol{p}}_{\boldsymbol{kl}}d^{\boldsymbol{k}}c^{\boldsymbol{l}}\overline{e}_{\boldsymbol{p}}\end{aligned} \tag{4.15}$$

$$\mathbf{1}<\boldsymbol{k}<\boldsymbol{n}\quad \mathbf{1}<\boldsymbol{l}<\boldsymbol{n}\quad \mathbf{1}<\boldsymbol{p}<\boldsymbol{n}$$

The equation (4.11) follows from the equation (4.15). □

Corollary 4.6. $\overline{e}_{\boldsymbol{k}}\,\overline{e}_{\boldsymbol{k}}\in\operatorname{Re}A$ □

Theorem 4.7. *(*$\boldsymbol{k}>0$, $\boldsymbol{l}>0$*)*

$$\overline{e}_{\boldsymbol{k}}\,\overline{e}_{\boldsymbol{l}}=(\overline{e}_{\boldsymbol{l}}\,\overline{e}_{\boldsymbol{k}})^* \tag{4.16}$$

Proof. From the equation (4.11), it follows that ($\boldsymbol{p}=\mathbf{1}$, ..., $\boldsymbol{n}$)

$$\begin{aligned}\overline{e}_{\boldsymbol{k}}\,\overline{e}_{\boldsymbol{l}}&=C^{\mathbf{0}}_{\boldsymbol{ab}}e^{\boldsymbol{a}}_{\boldsymbol{k}}e^{\boldsymbol{b}}_{\boldsymbol{l}}\overline{e}_{\mathbf{0}}+C^{\boldsymbol{p}}_{\boldsymbol{ab}}e^{\boldsymbol{a}}_{\boldsymbol{k}}e^{\boldsymbol{b}}_{\boldsymbol{l}}\overline{e}_{\boldsymbol{p}}\\&=C^{\mathbf{0}}_{\boldsymbol{ba}}e^{\boldsymbol{b}}_{\boldsymbol{l}}e^{\boldsymbol{a}}_{\boldsymbol{k}}\overline{e}_{\mathbf{0}}-C^{\boldsymbol{p}}_{\boldsymbol{ba}}e^{\boldsymbol{b}}_{\boldsymbol{l}}e^{\boldsymbol{a}}_{\boldsymbol{k}}\overline{e}_{\boldsymbol{p}}\\&=(C^{\mathbf{0}}_{\boldsymbol{ba}}e^{\boldsymbol{b}}_{\boldsymbol{l}}e^{\boldsymbol{a}}_{\boldsymbol{k}}\overline{e}_{\mathbf{0}}+C^{\boldsymbol{p}}_{\boldsymbol{ba}}e^{\boldsymbol{b}}_{\boldsymbol{l}}e^{\boldsymbol{a}}_{\boldsymbol{k}}\overline{e}_{\boldsymbol{p}})^*\\&=(\overline{e}_{\boldsymbol{l}}\,\overline{e}_{\boldsymbol{k}})^*\end{aligned} \tag{4.17}$$

The equation (4.16) follows from the equation (4.17). □

Theorem 4.8. *($k>0$, $l>0$)*

$$\overline{e}_k\,\overline{e}_l+\overline{e}_l\,\overline{e}_k\in\mathrm{Re}\,A \tag{4.18}$$

$$\overline{e}_k\,\overline{e}_l-\overline{e}_l\,\overline{e}_k\in\mathrm{Im}\,A \tag{4.19}$$

Proof. The theorem follows from the equation (4.16) and the theorem 4.4. □

Theorem 4.9. $dd^*\in\mathrm{Re}\,A$

Proof. From the equations (4.7), (4.11), it follows that $(p=1,...,n)$

$$\begin{aligned}dd^*&=(d^0\overline{e}_0+d^p\overline{e}_p)(d^0\overline{e}_0+d^q\overline{e}_q)^*\\&=(d^0\overline{e}_0+d^p\overline{e}_p)(d^0\overline{e}_0-d^q\overline{e}_q)\\&=d^0\overline{e}_0\,d^0\overline{e}_0-d^0\overline{e}_0\,d^q\overline{e}_q+d^p\overline{e}_p\,d^0\overline{e}_0-d^p\overline{e}_p\,d^q\overline{e}_q\\&=(d^0)^2\,\overline{e}_0-d^0d^p\overline{e}_p+d^0d^p\overline{e}_p-d^pd^q\overline{e}_p\,\overline{e}_q\end{aligned}\tag{4.20}$$

First 3 items in the expression (4.20) have form

$$(d^0)^2\,\overline{e}_0-d^0d^p\overline{e}_p+d^0d^p\overline{e}_p=(d^0)^2\,\overline{e}_0\in\mathrm{Re}\,A \tag{4.21}$$

The last item in the expression (4.20) has form

$$-d^pd^q\overline{e}_p\,\overline{e}_q=-(d^p)^2\,\overline{e}_p\,\overline{e}_p-\sum_{q>p}d^pd^q(\overline{e}_p\,\overline{e}_q+\overline{e}_q\,\overline{e}_p) \tag{4.22}$$

According to the corollary 4.6 and statement (4.18)

$$-d^pd^q\overline{e}_p\,\overline{e}_q\in\mathrm{Re}\,A \tag{4.23}$$

The theorem follows from the equation (4.20) and statements (4.21), (4.23). □

We can represent the conjugation using the matrix I

$$\begin{array}{llllll}d^*=I\circ d & I^0_k= & \delta^0_k & I^k_0= & \delta^k_0 & k=0,...,n\\ & I^m_k & =-\delta^m_k & I^k_m & =-\delta^k_m & m=1,...,n\end{array}\tag{4.24}$$

Example 4.10. The product in the set of complex numbers C is commutative. However, complex field has subfield R. The vector space $\mathrm{Im}\,C$ has dimension 1 and the basis $\overline{e}_1=i$. Accordingly

$$c^*=c^0-c^1i$$

□

Example 4.11. The division ring of quaternions H has subfield R. The vector space $\mathrm{Im}\,H$ has dimension 3 and the basis

$$\overline{e}_1=i\quad\overline{e}_2=j\quad\overline{e}_3=k$$

Accordingly

$$d^*=d^0-d^1i-d^2j-d^3k$$

□

5. Antilinear Mapping of Algebra with Conjugation

Definition 5.1. Let A be the algebra with conjugation. $(\mathrm{Re}\,A)$-linear mapping

$$f:A\rightarrow A$$

is called $A\star$**-antilinear**, if the mapping f satisfies to the equation

$$f(da)=f(a)d^*\tag{5.1}$$

□

Theorem 5.2. *$A\star$-antilinear mapping of the algebra with conjugation A has form*

$$f(c)=dc^*=dI\circ c\quad d\in A$$

Let $\overline{\overline{e}}$ be $(\mathrm{Re}\,A)$-basis of the algebra A. Coordinates of the mapping f

$$f(c)=c^l f_l^i\overline{e}_i\tag{5.2}$$

relative to the basis $\overline{\overline{e}}$ satisfy to the equation

$$f_p^j=f_0^k I_p^i C_{ki}^j\tag{5.3}$$

$$d=f_0^i\,\overline{e}_i\tag{5.4}$$

Proof. According to the definition 5.1, $A\star$-antilinear mapping satisfies to equations

$$f(ac)=(ac)^i f_i^j\overline{e}_j=a^p c^q C_{pq}^i f_i^j\overline{e}_j\tag{5.5}$$

$$f(c)a^*=(f(c))^k(a^*)^i C_{ki}^j\overline{e}_j=c^l f_l^k I_q^i a^q C_{ki}^j\overline{e}_j\tag{5.6}$$

From equations (5.5), (5.6), it follows that

$$a^p c^q C_{pq}^i f_i^j\overline{e}_j=c^q f_q^k I_p^i a^p C_{ki}^j\overline{e}_j\tag{5.7}$$

The equation

$$C_{pq}^i f_i^j=f_q^k I_p^i C_{ki}^j\tag{5.8}$$

follows from the equation (5.7).

From the equation (5.8) it follows that[6]

$$C_{p0}^i f_i^j=f_0^k I_p^i C_{ki}^j\tag{5.9}$$

From equations (3.5), (4.24), (5.9), it follows that

$$\delta_p^i f_i^j=f_0^k I_p^i C_{ki}^j\tag{5.10}$$

The equation (5.3) follows from the equation (5.10).

From equations (5.2), (5.3), it follows that

$$f(c)=c^l f_0^k I_l^i C_{ki}^j\overline{e}_j=f_0^k c^l I_l^i C_{ki}^j\overline{e}_j=f_0^k(c^*)^i C_{ki}^j\overline{e}_j\tag{5.11}$$

The equation (5.4) follows from the equations (3.6), (5.11). □

---

[6] Let $q=0$.

6. REFERENCES

[1] Serge Lang, Algebra, Springer, 2002

[2] Aleks Kleyn, Introduction into Calculus over Division Ring,
eprint arXiv:0812.4763 (2010)

[3] Aleks Kleyn, Quaternion Rhapsody,
eprint arXiv:0909.0855 (2010)

[4] Aleks Kleyn, Linear Mappings of Free Algebra: First Steps in Noncommutative Linear Algebra,
Lambert Academic Publishing, 2010

[5] Richard D. Schafer, An Introduction to Nonassociative Algebras, Dover Publications, Inc., New York, 1995

7. INDEX

algebra with conjugation 5
$A\star$-antilinear mapping of algebra with conjugation 8

conjugation in algebra 5

scalar algebra of algebra 5
scalar of element of algebra 5

vector module of algebra 5
vector of element of algebra 5

8. Special Symbols and Notations

d^* conjugation in algebra 5

$\operatorname{Im} A$ vector module of algebra A 5
$\operatorname{Im} d$ vector of element d of algebra 4

$\operatorname{Re} A$ scalar algebra of algebra A 5
$\operatorname{Re} d$ scalar of element d of algebra 4

arXiv:1105.4307v2 [math.RA] 9 Apr 2012

Алгебра с сопряжением

Александр Клейн

Аннотация. В статье рассмотрены свойства и отображения свободной алгебры с единицей. Рассмотренно также отображение сопряжения свободной алгебры с единицей.

Содержание

1. Предисловие . . . 1
2. Соглашения . . . 2
3. Алгебра с единицей . . . 2
4. Алгебра с сопряжением . . . 5
5. Антилинейное отображение алгебры с сопряжением . . . 8
6. Список литературы . . . 9
7. Предметный указатель . . . 10
8. Специальные символы и обозначения . . . 11

1. Предисловие

Когда я начал писать книгу [2], я понял, что классическое определение линейного отображения

$$y = ax \quad y = xa$$

ограничивает наши возможности изучать производную отображения в тело. Интуитивно было ясно, что линейное отображение в тело можно представить в виде суммы слагаемых вида

$$axb$$

Последующее исследование структуры линейного отображения в тело было завершено в книге [4]. При этом я расширил круг рассматриваемых алгебр, перейдя к рассмотрению свободных алгебр над коммутативным кольцом.

Если множество A является D-алгеброй, то на множестве A определено несколько алгебраических структур. Согласно определению [4]-2.2.1, A является D-модулем. Если мы ограничимся рассмотрением операций сложения и

Aleks_Kleyn@MailAPS.org.
http://sites.google.com/site/AleksKleyn/.
http://arxiv.org/a/kleyn_a_1.
http://AleksKleyn.blogspot.com/.

умножения в D-алгебре A, то мы можем рассматривать D-алгебру A как кольцо (возможно, неассоциативное). Соответственно этому мы можем рассматривать D-алгебру A как $A\star$-модуль размерности 1. Поэтому важно изучать отображения, сохраняющих ту или иную алгебраическую структуру на множестве A.

Эта статья рассматривает $A\star$-линейные отображения $D\star$-алгебры A.

Концепция алгебры с сопряжением естественно вписывается в теорию D-алгебры с единицей. Однако я полагаю, что отображение сопряжения можно определить также в случае, если алгебра не имеет единицу. Однако ответ на этот вопрос требует дополнительного исследования.

2. Соглашения

Соглашение 2.1. *Пусть A - свободная конечно мерная алгебра. При разложении элемента алгебры A относительно базиса $\overline{\overline{e}}$ мы пользуемся одной и той же корневой буквой для обозначения этого элемента и его координат. Однако в алгебре не принято использовать векторные обозначения. В выражении a^2 не ясно - это компонента разложения элемента a относительно базиса или это операция возведения в степень. Для облегчения чтения текста мы будем индекс элемента алгебры выделять цветом. Например,*

$$a=a^{i}\overline{e}_{i}$$

$\square$

Соглашение 2.2. *Если свободная конечномерная алгебра имеет единицу, то мы будем отождествлять вектор базиса $\overline{e}_{0}$ с единицей алгебры.* $\square$

Соглашение 2.3. *Хотя алгебра является свободным модулем над некоторым кольцом, мы не пользуемся векторными обозначениями при записи элементов алгебры. В тех случаях, когда я рассматриваю матрицу координат элемента алгебры, я буду пользоваться векторными обозначениями для записи соответствующего элемента. Чтобы не возникала неоднозначность при записи сопряжения, я буду обозначать a^* элемент, сопряжённый элементу a.* $\square$

Без сомнения, у читателя могут быть вопросы, замечания, возражения. Я буду признателен любому отзыву.

3. Алгебра с единицей

Пусть D - коммутативное кольцо. Пусть A - D-алгебра с единицей 1. Следовательно, определено эффективное представление

$$f_{1,2}:d\circ a=da\quad d\in D\quad a\in A \tag{3.1}$$

кольца D в алгебре A.

Теорема 3.1. *Пусть D - коммутативное кольцо. Пусть A - D-алгебра. Пусть*

$$f:A^n\to A$$

n-линейное отображение в D-алгебру A. Пусть

$$g:A^m\to A$$

m-линейное отображение в D-алгебру A. Тогда если для заданного i в выражении

$$f\circ(x_1,...,x_n)$$

выполнить замену

$$x_i=g\circ(y_1,...,y_m)$$

то полученное выражение

$$h(x_1,...,\widehat{x_i},...,x_n,y_1,...,y_m)=f\circ(x_1,...,g\circ(y_1,...,y_m),...,x_n)$$

является $(m+n-1)$-линейным отображением.

Доказательство. Поскольку полилинейное отображение линейно по каждому аргументу, то отображение h линейно по x^j, $j\neq i$. Так как отображение f линейно по x^i и для любого k отображение g линейно по y^k, то согласно теореме [4]-2.4.1 отображение h линейно по y^k. □

Теорема 3.2. *Пусть D - коммутативное кольцо. Пусть A - D-алгебра. Тогда*

$$d(ab)=(da)b=a(db)\quad d\in D\quad a,b\in A \tag{3.2}$$

Доказательство. Пусть f - билинейное отображение, соответствующее операции произведения в D-алгебре A.[1] Тогда

$$df(ab)=f(da,b)=f(a,db)\quad d\in D\quad a,b\in A \tag{3.3}$$

Равенство (3.2) следует из равенства (3.3). □

Теорема 3.3. *Пусть D - коммутативное кольцо.*[2] *Коммутатор D-алгебры A - билинейное отображение. Ассоциатор D-алгебры A - 3-линейное отображение.*

Доказательство. Согласно определениям [4]-2.2.1, [4]-2.2.4, коммутатор является билинейным отображением. Согласно определениям [4]-2.2.1, [4]-2.2.5 и теореме 3.1, ассоциатор является 3-линейным отображением. □

Теорема 3.4. *Пусть D - коммутативное кольцо. Пусть A - D-алгебра с единицей e. Тогда мы можем отождествить $d\in D$ и $d1\in A$. При этом $D\subseteq Z(A)$.*[3]

Доказательство. Из теоремы 3.3 следует

$$\begin{aligned}(d,a,b)&=d(1,a,b)=d((1a)b-1(ab))=d(ab-ab)=0\\(a,d,b)&=d(a,1,b)=d((a1)b-a(1b))=d(ab-ab)=0\\(a,b,d)&=d(a,b,1)=d((ab)1-a(b1))=d(ab-ab)=0\end{aligned}$$

Следовательно, $D\subseteq N(A)$. Если в равенстве (3.2) мы положим $b=1$, то мы получим

$$da=ad\quad d\in D\quad a\in A \tag{3.4}$$

---

[1] Я рассматриваю определение произведения в D-алгебре согласно определению [4]-2.2.1.

[2] Определение коммутатора дано в определении [4]-2.2.4. Определение ассоциатора дано в определении [4]-2.2.5. Смотри также [5], с. 13.

[3] Определение ядра $N(A)$ алгебры A дано в определении [4]-2.2.7; определение центра $Z(A)$ алгебры A дано в определении [4]-2.2.8; смотри также [5], с. 13, 14.

Из равенства (3.4) следует, что коммутатор имеет вид

$$[d,a]=0$$

□

Теорема 3.5. *Структурные константы D-алгебры с единицей e удовлетворяют условию*

$$C^l_{0k}=C^l_{k0}=\delta^k_l \tag{3.5}$$

Доказательство. Равенство (3.5) является следствием равенства $1d=d1=d$. □

Теорема 3.6. *$A\star$-линейное отображение D-алгебры A с единицей имеет вид*

$$f(c)=cd \quad d\in A \tag{3.6}$$

Пусть $\overline{\overline{e}}$ - $(\mathrm{Re}\,A)$-базис алгебры A. Координаты отображения f

$$f(c)=c^l f^i_l\overline{e}_i \tag{3.7}$$

относительно базиса $\overline{\overline{e}}$ удовлетворяют равенству

$$f^j_k=f^i_0C^j_{ki} \tag{3.8}$$

$$d=f^i_0\overline{e}_i \tag{3.9}$$

Доказательство. $A\star$-линейное отображение удовлетворяет равенствам

$$f(ac)=(ac)^i f^j_i\overline{e}_j=a^kc^lC^i_{kl}f^j_i\overline{e}_j \tag{3.10}$$

$$af(c)=a^k(f(c))^iC^j_{ki}\overline{e}_j=a^kc^lf^i_lC^j_{ki}\overline{e}_j \tag{3.11}$$

Из равенств (3.10), (3.11), следует

$$a^kc^lC^i_{kl}f^j_i\overline{e}_j=a^kc^lf^i_lC^j_{ki}\overline{e}_j \tag{3.12}$$

Равенство

$$C^i_{kl}f^j_i=f^i_lC^j_{ki} \tag{3.13}$$

следует из равенства (3.12).

Из равенства (3.13) следует[4]

$$C^i_{k0}f^j_i=f^i_0C^j_{ki} \tag{3.14}$$

Из равенств (3.5), (3.14), следует

$$\delta^i_kf^j_i=f^i_0C^j_{ki} \tag{3.15}$$

Равенство (3.8) следует из равенства (3.15).

Из равенств (3.7), (3.8) следует

$$f(c)=c^lf^j_0C^i_{lj}\overline{e}_i \tag{3.16}$$

Равенство (3.9) следует из равенств (3.6), (3.16). □

---

[4] Пусть $l=0$.

4. АЛГЕБРА С СОПРЯЖЕНИЕМ

Пусть D - коммутативное кольцо. Пусть A - D-алгебра с единицей e, $A\neq D$. Согласно теореме 3.4, мы можем отождествить кольцо D с подалгеброй D-алгебры A. Следовательно, D-алгебра A имеет нетривиальный центр $Z(A)$.

Пусть существует подалгебра F алгебры A такая, что $F\neq A$, $D\subseteq F\subseteq Z(A)$, и алгебра A является свободным модулем над кольцом F. Пусть $\overline{\overline{e}}$ - базис свободного модуля A над кольцом F. Мы будем полагать $\overline{e}_0=1$.

Рассмотрим отображения

$$\begin{aligned}\mathrm{Re}\,&: A\to A\\ \mathrm{Im}\,&: A\to A\end{aligned}$$

определенные равенством

$$\mathrm{Re}\,d=d^0\qquad \mathrm{Im}\,d=d-d^0\qquad d\in D\qquad d=d^i\overline{e}_i \tag{4.1}$$

Выражение $\mathrm{Re}\,d$ называется **скаляром элемента** d. Выражение $\mathrm{Im}\,d$ называется **вектором элемента** d.[5]

Согласно (4.1)

$$F=\{d\in A:\mathrm{Re}\,d=d\}$$

Мы будем пользоваться записью $\mathrm{Re}\,A$ для обозначения **алгебры скаляров алгебры** A.

Теорема 4.1. *Множество*

$$\mathrm{Im}\,A=\{d\in A:\mathrm{Re}\,d=0\} \tag{4.2}$$

является $(\mathrm{Re}\,A)$*-модулем, который мы называем* **модуль векторов алгебры** A.

$$A=\mathrm{Re}\,A\oplus\mathrm{Im}\,A \tag{4.3}$$

Доказательство. Пусть c, $d\in\mathrm{Im}\,A$. Тогда $c_0=d_0=0$. Следовательно,

$$(c+d)_0=c_0+d_0=0$$

Если $a\in\mathrm{Re}\,A$, то

$$(ad)_0=ad_0=0$$

Следовательно, $\mathrm{Im}\,A$ является $(\mathrm{Re}\,A)$-модулем.

Последовательность модулей

$$\begin{tikzcd}0 \arrow[r] & \mathrm{Re}\,A \arrow[r, "\mathrm{id}"] & A \arrow[r, "\mathrm{Im}"] & \mathrm{Im}\,A \arrow[r] & 0\end{tikzcd}$$

является точной последовательностью. Согласно определению (4.1) отображения Re, следующая диаграмма коммутативна

$$\begin{tikzcd}\mathrm{Re}\,A \arrow[r, "\mathrm{id}"] \arrow[d, "\mathrm{id}"'] & A \arrow[ld, "\mathrm{Re}"] \\ \mathrm{Re}\,A & \end{tikzcd}$$

Согласно утверждению (2) предложения [1]-III.3.3,

$$A=\mathrm{id}(\mathrm{Re}\,A)\oplus\ker\mathrm{Re} \tag{4.4}$$

---

[5] В разделе [3]-7, я рассматриваю пример алгебры, в которой определены две операции сопряжения. Мы будем полагать, что операция сопряжения выбрана.

Согласно определению (4.2)

$$\ker \mathrm{Re} = \{d \in A : \mathrm{Re}\, d = 0\} = \mathrm{Im}\, A \tag{4.5}$$

Равенство (4.3) следует из равенств (4.4), (4.5). □

Согласно теореме 4.1, однозначно определенно представление

$$d = \mathrm{Re}\, d + \mathrm{Im}\, d \tag{4.6}$$

Определение 4.2. Отображение

$$d^* = \mathrm{Re}\, d - \mathrm{Im}\, d \tag{4.7}$$

называется **сопряжением в алгебре** при условии, если это отображение удовлетворяет равенству

$$(cd)^* = d^*\, c^* \tag{4.8}$$

$(\mathrm{Re}\, A)$-алгебра A, в которой определено сопряжение, называется **алгеброй с сопряжением**. □

Следствие 4.3. $(d^*)^* = d$ □

Теорема 4.4.

$$d + d^* \in \mathrm{Re}\, A \tag{4.9}$$

$$d - d^* \in \mathrm{Im}\, A \tag{4.10}$$

Доказательство. Утверждение теоремы следует из равенств (4.6), (4.7). □

Теорема 4.5. $(\mathrm{Re}\, A)$*-алгебра* A *является алгеброй с сопряжением, если структурные константы* $(\mathrm{Re}\, A)$*-алгебры* A *удовлетворяют условию*

$$C^{0}_{kl} = C^{0}_{lk} \quad C^{p}_{kl} = -C^{p}_{lk} \tag{4.11}$$

$$1 < k < n \quad 1 < l < n \quad 1 < p < n$$

Доказательство. Из равенств (4.6), (4.7) следует

$$\begin{aligned}(cd)^* &= (\mathrm{Re}\, c\ \mathrm{Re}\, d + \mathrm{Re}\, c\ \mathrm{Im}\, d + \mathrm{Im}\, c\ \mathrm{Re}\, d + \mathrm{Im}\, c\ \mathrm{Im}\, d)^*\\ &= \mathrm{Re}\, c\ \mathrm{Re}\, d - \mathrm{Re}\, c\ \mathrm{Im}\, d - \mathrm{Im}\, c\ \mathrm{Re}\, d + (\mathrm{Im}\, c\ \mathrm{Im}\, d)^*\end{aligned} \tag{4.12}$$

$$\begin{aligned}d^*\, c^* &= (\mathrm{Re}\, d - \mathrm{Im}\, d)(\mathrm{Re}\, c - \mathrm{Im}\, c)\\ &= \mathrm{Re}\, d\ \mathrm{Re}\, c - \mathrm{Re}\, d\ \mathrm{Im}\, c - \mathrm{Im}\, d\ \mathrm{Re}\, c + \mathrm{Im}\, d\ \mathrm{Im}\, c\end{aligned} \tag{4.13}$$

Из равенств (4.8), (4.12), (4.13), следует

$$(\mathrm{Im}\, c\ \mathrm{Im}\, d)^* = \mathrm{Im}\, d\ \mathrm{Im}\, c \tag{4.14}$$

Пусть $\overline{\overline{e}}$ - базис $(\mathrm{Re}\, D)$-алгебры D. Из равенства (4.14) следует

$$\begin{aligned}(C^{0}_{kl} c^{k} d^{l} + C^{p}_{kl} c^{k} d^{l} \overline{e}_{p})^* &= C^{0}_{kl} c^{k} d^{l} - C^{p}_{kl} c^{k} d^{l} \overline{e}_{p}\\ &= C^{0}_{kl} d^{k} c^{l} + C^{p}_{kl} d^{k} c^{l} \overline{e}_{p}\end{aligned} \tag{4.15}$$

$$1 < k < n \quad 1 < l < n \quad 1 < p < n$$

Равенство (4.11) следует из равенства (4.15). □

Следствие 4.6. $\overline{e}_k\,\overline{e}_k \in \mathrm{Re}\,A$ $\square$

Теорема 4.7. *($k>0$, $l>0$)*

$$\overline{e}_k\,\overline{e}_l = (\overline{e}_l\,\overline{e}_k)^* \tag{4.16}$$

Доказательство. Из равенства (4.11) следует $(p=1,\ ...,\ n)$

$$\begin{aligned}\overline{e}_k\,\overline{e}_l &= C^0_{ab}e^a_k e^b_l \overline{e}_0 + C^p_{ab}e^a_k e^b_l \overline{e}_p\\ &= C^0_{ba}e^b_l e^a_k \overline{e}_0 - C^p_{ba}e^b_l e^a_k \overline{e}_p\\ &= (C^0_{ba}e^b_l e^a_k \overline{e}_0 + C^p_{ba}e^b_l e^a_k \overline{e}_p)^*\\ &= (\overline{e}_l\,\overline{e}_k)^*\end{aligned} \tag{4.17}$$

Равенство (4.16) следует из равенства (4.17). $\square$

Теорема 4.8. *($k>0$, $l>0$)*

$$\overline{e}_k\,\overline{e}_l + \overline{e}_l\,\overline{e}_k \in \mathrm{Re}\,A \tag{4.18}$$

$$\overline{e}_k\,\overline{e}_l - \overline{e}_l\,\overline{e}_k \in \mathrm{Im}\,A \tag{4.19}$$

Доказательство. Утверждение теоремы следует из равенства (4.16) и теоремы 4.4. $\square$

Теорема 4.9. $dd^* \in \mathrm{Re}\,A$

Доказательство. Из равенств (4.7), (4.11) следует $(p=1,\ ...,\ n)$

$$\begin{aligned}dd^* &= (d^0\overline{e}_0 + d^p\overline{e}_p)(d^0\overline{e}_0 + d^q\overline{e}_q)^*\\ &= (d^0\overline{e}_0 + d^p\overline{e}_p)(d^0\overline{e}_0 - d^q\overline{e}_q)\\ &= d^0\overline{e}_0\,d^0\overline{e}_0 - d^0\overline{e}_0\,d^q\overline{e}_q + d^p\overline{e}_p\,d^0\overline{e}_0 - d^p\overline{e}_p\,d^q\overline{e}_q\\ &= (d^0)^2\,\overline{e}_0 - d^0d^p\overline{e}_p + d^0d^p\overline{e}_p - d^pd^q\overline{e}_p\,\overline{e}_q\end{aligned} \tag{4.20}$$

Первые 3 слагаемых в выражении (4.20) имеют вид

$$(d^0)^2\,\overline{e}_0 - d^0d^p\overline{e}_p + d^0d^p\overline{e}_p = (d^0)^2\,\overline{e}_0 \in \mathrm{Re}\,A \tag{4.21}$$

Последнее слагаемое в выражении (4.20) имеет вид

$$-\,d^pd^q\overline{e}_p\,\overline{e}_q = -(d^p)^2\,\overline{e}_p\,\overline{e}_p - \sum_{q>p} d^pd^q(\overline{e}_p\,\overline{e}_q + \overline{e}_q\,\overline{e}_p) \tag{4.22}$$

Согласно следствию 4.6 и утверждению (4.18)

$$-\,d^pd^q\overline{e}_p\,\overline{e}_q \in \mathrm{Re}\,A \tag{4.23}$$

Утверждение теоремы следует из равенства (4.20) и утверждений (4.21), (4.23). $\square$

Отображение сопряжения можно представить с помощью матрицы I

$$\begin{array}{llll} d^* = I\circ d & I^0_k = \delta^0_k & I^k_0 = \delta^k_0 & k=0,...,n\\ & I^m_k = -\,\delta^m_k & I^k_m = -\,\delta^k_m & m=1,...,n\end{array} \tag{4.24}$$

Пример 4.10. Произведение в множестве комплексных чисел C коммутативно. Однако поле комлексных чисел содержит подполе R. Векторное пространство $\mathrm{Im}\,C$ имеет размерность 1 и базис $\overline{e}_1 = i$. Соответственно

$$c^* = c^0 - c^1 i$$

□

Пример 4.11. Тело кватернионов H содержит подполе R. Векторное пространство $\mathrm{Im}\,H$ имеет размерность 3 и базис

$$\overline{e}_1 = i \quad \overline{e}_2 = j \quad \overline{e}_3 = k$$

Соответственно

$$d^* = d^0 - d^1 i - d^2 j - d^3 k$$

□

5. АНТИЛИНЕЙНОЕ ОТОБРАЖЕНИЕ АЛГЕБРЫ С СОПРЯЖЕНИЕМ

Определение 5.1. Пусть A - алгебра с сопряжением. $(\mathrm{Re}\,A)$-линейное отображение

$$f : A \to A$$

называется $A\star$**-антилинейным** если отображение f удовлетворяет равенству

$$f(da) = f(a)d^* \tag{5.1}$$

□

Теорема 5.2. *$A\star$-антилинейное отображение алгебры с сопряжением A имеет вид*

$$f(c) = dc^* = dI \circ c \quad d \in A$$

Пусть $\overline{\overline{e}}$ - $(\mathrm{Re}\,A)$-базис алгебры A. Координаты отображения f

$$f(c) = c^l f^i_l \overline{e}_i \tag{5.2}$$

относительно базиса $\overline{\overline{e}}$ удовлетворяют равенству

$$f^j_p = f^k_0 I^i_p C^j_{ki} \tag{5.3}$$

$$d = f^i_0 \overline{e}_i \tag{5.4}$$

Доказательство. Согласно определению 5.1, $A\star$-антилинейное отображение удовлетворяет равенствам

$$f(ac) = (ac)^i f^j_i \overline{e}_j = a^p c^q C^i_{pq} f^j_i \overline{e}_j \tag{5.5}$$

$$f(c)a^* = (f(c))^k (a^*)^i C^j_{ki} \overline{e}_j = c^l f^k_l I^i_q a^q C^j_{ki} \overline{e}_j \tag{5.6}$$

Из равенств (5.5), (5.6), следует

$$a^p c^q C^i_{pq} f^j_i \overline{e}_j = c^q f^k_q I^i_p a^p C^j_{ki} \overline{e}_j \tag{5.7}$$

Равенство

$$C^i_{pq} f^j_i = f^k_q I^i_p C^j_{ki} \tag{5.8}$$

следует из равенства (5.7).

Из равенства (5.8) следует[6]

$$C^{i}_{p0}f^{j}_{i}=f^{k}_{0}I^{i}_{p}C^{j}_{ki} \tag{5.9}$$

Из равенств (3.5), (4.24), (5.9), следует

$$\delta^{i}_{p}f^{j}_{i}=f^{k}_{0}I^{i}_{p}C^{j}_{ki} \tag{5.10}$$

Равенство (5.3) следует из равенства (5.10).

Из равенств (5.2), (5.3) следует

$$f(c)=c^{l}f^{k}_{0}I^{i}_{l}C^{j}_{ki}\overline{e}_{j}=f^{k}_{0}c^{l}I^{i}_{l}C^{j}_{ki}\overline{e}_{j}=f^{k}_{0}(c^{*})^{i}C^{j}_{ki}\overline{e}_{j} \tag{5.11}$$

Равенство (5.4) следует из равенств (3.6), (5.11). □

6. Список литературы

[1] Серж Ленг, Алгебра, М. Мир, 1968

[2] Александр Клейн, Введение в математический анализ над телом,
eprint arXiv:0812.4763 (2010)

[3] Александр Клейн, Этюд о кватернионах,
eprint arXiv:0909.0855 (2010)

[4] Aleks Kleyn, Linear Mappings of Free Algebra: First Steps in Noncommutative Linear Algebra,
Lambert Academic Publishing, 2010

[5] Richard D. Schafer, An Introduction to Nonassociative Algebras, Dover Publications, Inc., New York, 1995

---

[6] Пусть $q=0$.

7. Предметный указатель

$A\star$-антилинейное отображение алгебры с сопряжением 8

алгебра с сопряжением 6
алгебра скаляров алгебры 5

вектор элемента алгебры 5

модуль векторов алгебры 5

скаляр элемента алгебры 5
сопряжение в алгебре 6

8. Специальные символы и обозначения

d^* сопряжение в алгебре 6

$\operatorname{Im} A$ модуль векторов алгебры A 5
$\operatorname{Im} d$ вектор элемента d алгебры 5

$\operatorname{Re} A$ алгебра скаляров алгебры A 5
$\operatorname{Re} d$ скаляр элемента d алгебры 5